

Figure I.C.4.a: Diabetes Mellitus (DM)



When provider documentation refers to use, abuse, and dependence of the same substance (e.g., alcohol, opioid, cannabis, etc.), only one code should be assigned to identify the pattern of use based on the highest step. Example: The provider documents tobacco use and abuse. Report only tobacco abuse. If alcohol use and dependence is documented, report alcohol dependence.

© AAPC. All rights reserved.

Figure I.C.5.b.2: Psychoactive Substance Use, Abuse and Dependence

Chapter 5: Mental, Behavioral and Neurodevelopmental disorders (F01–F99)

a. Pain disorders related to psychological factors

Assign code F45.41, for pain that is exclusively related to psychological disorders. As indicated by the Excludes 1 note under category G89, a code from category G89 should not be assigned with code F45.41.

Code F45.42, Pain disorders with related psychological factors, should be used with a code from category G89, Pain, not elsewhere classified, if there is documentation of a psychological component for a patient with acute or chronic pain.

See Section I.C.6. Pain

b. Mental and behavioral disorders due to psychoactive substance use

1) In Remission

Selection of codes for "in remission" for categories F10-F19, Mental and behavioral disorders due to psychoactive substance use (categories F10-F19 with -.11, -.21) requires the provider's clinical judgment. The appropriate codes for "in remission" are assigned only